



## **Historic, Archive Document**

Do not assume content reflects current scientific knowledge, policies, or practices.



1  
P6985d

S. D.—83. [This leaflet is distributed only with the seeds to which it relates.]

## United States Department of Agriculture,

### BUREAU OF PLANT INDUSTRY,

New and Rare Seed Distribution,

WASHINGTON, D. C.

## ALFALFA.

Instructions adapted to the New England States and New York.

**OBJECT OF THE DISTRIBUTION.**—The distribution of new and rare seeds has for its object the dissemination of new and rare crops, improved strains of staple crops, and high-grade seed of crops new to sections where the data of the department indicate such crops to be of considerable promise. Each package contains a sufficient quantity for a preliminary trial, and where it is at all practicable the recipient is urged to use the seed for the production of stocks for future plantings. It is believed that if this practice is followed consistently it will result in a material improvement in the crops of the country.

Please make a full report on the inclosed blank regarding the results you obtain with the seed.

### DESCRIPTION.

Alfalfa (*Medicago sativa*) is a deep-rooted, hardy, perennial forage plant belonging to the family which includes beans, peas, and clover. It occupies the same place in western agriculture that clover fills in the Northeastern States. In chemical composition it resembles the clovers, peas, and allied legumes in having a high protein content, but it slightly outranks any of these which are now in cultivation in both total and digestible protein. Good clover hay is almost equal to alfalfa in feeding value, but the number of crops alfalfa produces in one season makes the total yield per acre greater than that of clover. Since it is perennial, it will last a number of years unless crowded out by weeds or otherwise destroyed.

### SOIL REQUIREMENTS.

A deep, fertile, well-drained, nonacid soil reasonably free from weeds is required. It is practically useless to sow alfalfa on thin soils where the bed-rock approaches the surface, on land underlain with hardpan, or in locations where the subsoil is so compact that the roots can not penetrate it to considerable depths. It is also equally useless to attempt to grow alfalfa on land where the water table comes near the surface. For the purpose of ascertaining the character of the soil and subsoil and also the depth to the water table, frequent borings should be made with a soil auger. In determining the adaptability of a tract of land to alfalfa this instrument will generally be of greater assistance than a chemical analysis of the soil.

Not only should the land have good underdrainage, but the surface should have sufficient slope to carry off the surplus water readily. In fields that are too level, or in pockets, the formation of ice on the surface is fatal to alfalfa.



In this climate ice may form on the surface even on considerable slopes, but this is a danger that can not be avoided and is least on these slopes. Rich river or creek bottom lands that are well drained are admirably suited to the crop. On such lands overflows may do no serious damage, provided they are not of long duration and come at a period of the year when the growth is practically dormant.

Limestone soils are in general especially well suited to the production of alfalfa, but even these soils are frequently acid and require liming before they will grow the crop successfully.

#### PRECEDING CROP.

When once started under favorable soil conditions, weeds, including bluegrass, are likely to prove the most dangerous enemy. For this reason it is best to precede the alfalfa for one or two years with a clean cultivated crop, such as truck, potatoes, or corn. Summer fallowing also offers an excellent opportunity for eradicating weeds, but this practice is objected to on the ground that it results in the loss of the use of the land for a season. Successful stands are also secured following wheat, oats, and rye, provided the land has previously been treated in such a manner as to destroy the weeds. In this section, however, some difficulty is encountered in getting these crops off in time to permit the proper preparation of the land for sowing alfalfa the same season. Generally speaking, sowing on sod land is not recommended.

#### PREPARATION OF LAND.

The tender nature of the alfalfa plants requires that the soil be in excellent tilth at the time of planting. Many of the failures to secure a good stand may be traced directly to the improper condition of the seed bed. The soil should be fine and loose for the surface 2 or 3 inches, and below that it should be sufficiently firm to favor capillary action, yet porous enough to insure good drainage and aeration. Such a condition can best be secured by plowing the land the previous fall. In the spring, as soon as the weeds put in an appearance, the land should be disked to destroy these weeds and further pulverize and settle the seed bed. From this time until the time of sowing, the land should be cultivated every 10 to 12 days, or at least with sufficient frequency to check the weed growth. If it is impracticable to plow in the fall, the land should be plowed at least six weeks before sowing and harrowed frequently, in order that the ground may be in the proper condition at sowing time. Land on which small grain has been grown may be prepared by double-disking as soon as the grain is removed and then harrowing at frequent intervals until the seed is sown.

#### LIMING.

Practically all the soils in the region under consideration are benefited by applications of lime. The lime may be applied with a manure spreader, a fertilizer distributor, a lime distributor, or by hand. Any method which spreads the lime uniformly and at low cost is satisfactory. It should be applied two or three weeks before sowing in order that it may become thoroughly incorporated with the soil. At least a ton of burned lime is generally required, and larger applications are frequently necessary. If ground limestone or ground oyster shell is to be used, the quantity should be double that of burned lime. Experiments have shown very little difference in the final results obtained from the different forms of lime. Burned lime will give quicker results, but the ground limestone and ground oyster shell will finally give



the same benefit. The essential element in lime in any form is the calcium oxid, and it is recommended that the farmer use whichever form of lime is cheapest, based upon the percentage of this element present. Where the consumer pays the freight, it should be remembered that he will not only have to pay such charges on practically twice as much of the ground limestone as of the burned lime but will also be to the additional expense of hauling and spreading 2 tons of the former to 1 of the latter in order to obtain the same results.

#### FERTILIZING.

Well-rotted stable manure which is comparatively free from weed seeds is generally the most satisfactory fertilizer for alfalfa. Where the land is plowed in the fall it may be spread before plowing or it may be applied as a top-dressing during the early winter months. Beneficial results also follow heavy applications to the preceding crop.

Where stable manure is not available, a liberal application of commercial fertilizers, rich in phosphoric acid, should be made. The percentage of nitrogen may be low, but some nitrogen should be supplied for the young plants before they become inoculated and are able to secure their supply from the air. A combination which has given good results consists of 100 to 150 pounds of muriate of potash, 350 to 500 pounds of acid phosphate, and 50 to 75 pounds of nitrate of soda per acre. The cheapest and most satisfactory method for the consumer to obtain the desired combination is to purchase the various ingredients and mix them in the proper proportion.

#### INOCULATION.

Nitrogen-fixing bacteria should be provided unless the soil is known to be naturally supplied with these germs. This may be accomplished best by scattering over the area to be sown soil from a field upon which the crop has previously been successfully grown. The soil should be broadcast at the rate of 250 to 500 pounds per acre and harrowed as soon as practicable. Soil from the roots of sweet-clover plants also will inoculate alfalfa. Inoculation may also be accomplished by the use of an artificial culture, a limited quantity of which can be procured from the United States Department of Agriculture free of charge. Full instructions for use accompany each bottle of culture. The combined use of the soil and the artificial culture is recommended where both can be readily obtained.

#### SOWING.

The seed should be sown at the rate of 25 to 30 pounds to the acre, the heavier sowing being preferable, as it makes hay of a finer quality and helps to keep down the weeds. It may be drilled or sown broadcast with a wheelbarrow seeder and covered lightly with a smoothing harrow or weeder, care being taken to avoid covering deeper than 1 inch to 1½ inches. As a rule, a higher percentage of germination is secured from drilling, and such being the case the quantity of seed used per acre may be slightly decreased. A much more even stand can usually be secured by dividing the seed and sowing one-half each way of the field. The most successful stands of alfalfa are secured from sowing late in June or as soon thereafter as moisture conditions are favorable. It is seldom safe to sow much later than August 15, as the plants will not have sufficient time to become thoroughly established before winter sets in. Successful stands are occasionally secured from early spring sowing, although they are not generally recommended. A nurse crop should not be used unless the seed is sown in the spring, when 1½ bushels of oats or beardless barley, preferably the



latter, may be used to help keep down the weeds. This should be cut for hay; otherwise, the growth of the alfalfa may be seriously checked. Successful stands have also been secured from sowing with early canning peas. In some cases the sowing of alfalfa in standing corn in the latter part of July has given successful results, but this method can be expected to succeed only when conditions are highly favorable.

#### TREATMENT OF THE STAND.

In spring sowings, unless the weeds threaten to choke out the young plants, the alfalfa should not be clipped until it is from 12 to 15 inches high and in bloom. The cutter bar of the mower should be set high, as the young alfalfa is likely to be injured if cut low. If there is sufficient growth to be gathered with a rake it should be removed from the field. A fall sowing should not be mowed until the following spring, when it can be cut for hay each time the plants are pretty well in bloom or the basal shoots have made a good start. When the plants turn yellow, the crop should be cut immediately and removed from the field. If the stand becomes thin or patchy, the field should be plowed and reseeded. Attempts at patching up poor stands have generally proved futile. In cases where the soil has become compact from pasturing or where weeds are so numerous as to threaten considerable injury to the crop, cultivation in the spring or immediately after cutting with a spring-tooth harrow or "alfalfa harrow" may be advisable. No implements which will seriously injure the crowns and permit diseases to enter should be used. A top-dressing of well-rotted, weed-free stable manure, applied in the late fall or in winter, will not only furnish some protection but will also in most cases increase the yields the following season. If the stable manure is not available, a top-dressing of 300 to 500 pounds of acid phosphate per acre, with a small quantity of potash, may prove beneficial. Alfalfa makes good pasture for nearly all kinds of farm animals, but under no circumstances should it be pastured until it has become thoroughly established or when the ground is wet or frozen. It should not be pastured closely at any time, for the grazing down of the crowns will often result in its destruction. Owing to the difficulty of procuring a good stand in the Eastern States, it is very doubtful whether a farmer should take the chance or injuring a well-established stand by pasturing it at all.

#### TWO LEADING COMMERCIAL VARIETIES OF ALFALFA.

*Common alfalfa.*—Under this head is included the greater part of the alfalfa grown in the United States, the seed from the various sources frequently being designated by the State in which it was produced. Where alfalfa has been grown under a certain set of conditions for a considerable time, there is a tendency, through elimination, to produce a different type, presumably best adapted to the conditions under which it was developed. Thus, for instance, seed from fields that have been grown for several seed generations in Montana and the Dakotas may be somewhat more hardy than that grown farther south. Likewise, seed grown under semiarid conditions without irrigation may be superior for dry-land farming. In sections where winterkilling is not a factor, the ordinary types are recommended in preference to the so-called hardy alfalfa, as they generally produce somewhat heavier yields. An effort should be made to procure seed produced in approximately the same latitude as that in which it is to be sown.

*Grimm alfalfa.*—Grimm alfalfa was introduced into this country in 1857 from Baden, Germany, by Wendelin Grimm, of Carver County, Minn. Careful investigations indicate that it owes its superior hardiness to the fact that it is

the result of a natural cross between the common variety and the yellow-flowered alfalfa (*Medicago falcata*) and that by virtue of its being exposed to numerous severe winters the weaker plants were eliminated, leaving only the hardy ones to perpetuate the strain. Grimm alfalfa does not differ materially in appearance from the ordinary strain, so that the casual observer has difficulty in distinguishing one from the other. While a large percentage of its flowers are of the same color as those of common alfalfa, there are some that represent many shades of violet, yellow, and other hues. The taproots show a tendency to branch, and the crowns are inclined to be low set and spreading, characteristics which undoubtedly are of great importance in rendering the variety resistant to drought. Grimm alfalfa is one of the hardiest, if not the most hardy, of our commercial strains. It is recommended for sections where the winters are especially severe and where little protection is given by snow. In sections where winterkilling is not an important consideration it is not thought to be materially superior to common alfalfa.

#### PUBLICATIONS AVAILABLE.

For further details regarding alfalfa, see Farmers' Bulletins Nos. 757, Commercial Varieties of Alfalfa; 1229, Utilization of Alfalfa; and 1283, How to Grow Alfalfa, which will be sent free of charge upon application to the Secretary of Agriculture, Washington, D. C.

OCTOBER 30, 1923.









